Food Allergy Action Plan

Student's Name:		D.O.B:Teacher:			Place	
ALLERGY TO:					Child's Picture	
Asthmatic Yes* No *Higher risk for severe reaction					Here	
♦ STEP 1: TREATMENT ♦						
Symptoms:			Give Checked Medication**: To be determined		determined	
		allergen has been ingested, but no symptoms:	☐ EpiPen	☐ Antihistam	authorizing	
•	Mouth	Itching, tingling, or swelling of lips, tongue, mouth	☐ EpiPen	☐ Antihistamine		
	Skin	Hives, itchy rash, swelling of the face or extremities	☐ EpiPen	☐ Antihistamine		
=	Gut	Nausea, abdominal cramps, vomiting, diarrhea		☐ Antihistam	nine	
	Throat +	Tightening of throat, hoarseness, hacking cough	☐ EpiPen	☐ Antihistamine		
н	Lung †	Shortness of breath, repetitive coughing, wheezing	☐ EpiPen	☐ Antihistamine		
	Heart +	Thready pulse, low blood pressure, fainting, pale, blueness	☐ EpiPen	☐ Antihistamine		
	Other t		☐ EpiPen	☐ Antihistamine		
	 If reaction is progressing (several of the above areas affected), give 			☐ Antihistamine		
The severity of symptoms can quickly change. † Potentially life-threatening.						
DOSAGE Epinephrine: inject intramuscularly (circle one) EpiPen EpiPen Jr. (see reverse side for instructions)						
Antihistamine: give						
Other: give						
medication/dose/route						
◆ <u>STEP 2: EMERGENCY CALLS</u> ◆						
1. Call 911 (or Rescue Squad:). State that an allergic reaction has been treated, and additional epinephrine may be needed)						
2. Dr at						
3. Emergency contacts: Name/Relationship Phone Number(s)						
a 1.)		2.)				
EVEN IF PARENT/GUARDIAN CANNOT BE REACHED, DO NOT HESITATE TO MEDICATE OR TAKE CHILD TO MEDICAL FACILITY!						
Parent/Guardian Signature Date						
	tor's Signat			Date		